and they will be ſeparated by it, when held at the diſtance of three or four feet; let it be brought nearer, and they will ſtand farther apart; intirely withdraw it, and they will immediately come together. This experiment may be made with very ſmall braſs balls hung by ſilver wire; and will ſucceed as well with ſealing-wax made electrical, as with glaſs.

EXPERIMENT II.

If two cork-balls be ſuſpended by dry ſilk threads, the excited tube muſt be brought within eighteen inches before they will repel each other; which they will continue to do, for ſome time, after the tube is taken away.

As the balls in the firſt experiment are not inſulated, they cannot properly be ſaid to be electrified: but when they hang within the atmoſphere of the excited tube, they may attract and condenſe the electrical fluid round about them, and be ſeparated by the repulſion of its particles. It is conjectur'd alſo, that the balls at this time contain leſs than their common ſhare of the electrical fluid, on account of the repelling power of that which ſurrounds them; tho' ſome, perhaps, is continually entering and paſſing thro' the threads. And if that be the caſe, the reaſon is plain, why the balls hung by ſilk, in the ſecond experiment, muſt be in a much more denſe part of the atmoſphere of the tube, before they will repel each other. At the approach of an excited ſtick of wax to the balls, in the firſt experiment, the electrical fire is ſuppoſed to come through the threads into the balls, and be condenſed there, in its paſſage towards the wax: for, according to Mr *Franklin*, excited glaſs *emits* the electrical fluid, but excited wax *receives* it.

EXPERIMENT III.

Let a tin tube, of four or five feet in length, and about two inches inches in diameter, be inſulated by ſilk; and from one end of it let the cork-balls be ſuſpended by linen threads. Electrify it, by bringing the excited glaſs tube near the other end, ſo as that the balls may ſtand an inch and an half, or two inches apart: then, at the approach of the excited tube, they will by degrees loſe their repelling power, and come into contact; and as the tube is brought ſtill nearer, they will ſeparate again to as great a diſtance as before: in the return of the tube they will approach each other till they touch, and then repel as at firſt. If the tin-tube be electrified by wax, or the wire of a charg'd phial, the balls will be affected in the ſame manner at the approach of excited wax, or the wire of the phial.

EXPERIMENT IV.

Electrify the cork-balls as in the laſt experiment by glaſs; and at the approach of an excited ſtick of wax their repulſion will be increaſed. The effect will be the ſame, if the excited glaſs be brought towards them, when they have been electrified by wax.

The bringing the excited glaſs to the end, or edge of the

BENJAMIN FRANKLIN *Experiments and Observations on Electricity* LONDON, 1751 THE WARNOCK LIBRARY Octavo